


GLP-1 RAs AND CARDIOVASCULAR OUTCOMES IN TYPE 2 DIABETES



FDA guidance to industry on CVOT	ELIXA¹ (Lixisenatide) n=6068; follow-up 2.1 yrs.	LEADER¹ (Liraglutide) n=9340; duration 3.8 yrs.	SUSTAIN 6¹ (Semaglutide) n=3297; duration 2.1 yrs.	EXSCEL¹ (Exenatide ER) n=14752; follow-up 3.2 yrs.	REWIND¹ (Dulaglutide) n=9901; duration 5.4 yrs.	HARMONY¹ (Albiglutide) n=9463; duration 1.6 yrs.	PIONEER 6¹ (Oral semaglutide) n=3183; duration 1.3 yrs.	AMPLITUDE-O¹ (Efpeglenatide) n=4076; duration 1.8 yrs.	SOUL² (Oral semaglutide) n=9650; duration 5.2 yrs.
---	--	---	--	--	---	--	---	--	--

Baseline Characteristics and Use of Glucose-Lowering Agents Across Trials¹

Drug	ELIXA (n=6068)	LEADER (n=9340)	SUSTAIN 6 (n=3297)	EXSCEL (n=14752)	HARMONY OUTCOMES (n=9463)	REWIND (n=9901)	PIONEER-6 (n=3183)	AMPLITUDE-O (n=4076)
	Lixisenatide	Liraglutide	Semaglutide	Exenatide	Albiglutide	Dulaglutide	Semaglutide	Efpeglenatide
Administration route	Subcutaneous	Subcutaneous	Subcutaneous	Subcutaneous	Subcutaneous	Subcutaneous	Oral	Subcutaneous
Target dose	10 µg/d or 20 µg/d	1.8 mg/d	0.5 mg/wk or 1 mg/wk	2 mg/wk	30 mg/wk or 50 mg/wk	1.5 mg/wk	14 mg/d	4 mg/wk or 6 mg/wk
Age, yrs.	60±10	64±7	65±7	62±9	64±7	66±7	66±7	65±8
Female	31%	36%	39%	38%	31%	46%	32%	33%
Male	69%	64%	61%	62%	69%	54%	68%	67%
BMI kg/m ²	30.1±5.6	32.5±6.3	32.8±6.2	32.7±6.4	32.3±5.9	32.3±5.7	32.3±6.5	32.7±6.2
Diabetes duration, yrs.	9.2±8.2	12.8±8.0	13.9±8.1	13.1±8.3	14.2±8.8	10.5±7.2	14.9±8.5	15.4±8.8
HbA _{1c} %	7.7±1.3	8.7±1.6	8.7±1.5	8.1±1.0	8.7±1.5	7.3±1.1	8.2±1.6	8.9±1.5
Established cardiovascular disease	100%	81%	83%	73%	100%	31%	85%	90%
History of heart failure	22%	18%	24%	16%	20%	9%	12%	18%
Systolic blood pressure (mm Hg)	129±17	136±18	136±17	135±17	135±17	137±17	136±18	135±16
eGFR, mL/min per 1.73 m ² *	78±21	80 (NR)	80 (61–92)	77 (61–92)	79±25	77±23	74±21	72±22

Summary Results of Meta-Analysis for MACE and Its Components¹

	Main analysis with all 8 CVOTs (HR; I-squared)		Sensitivity analyses minus ELIXA (HR; I-squared)	
MACE	0.86 (0.80 to 0.93)	◆	0.85 (0.80 to 0.90)	◆
CV death	0.87 (0.80 to 0.94)	◆	0.85 (0.78 to 0.93)	◆
Fatal or non fatal MI	0.90 (0.83 to 0.98)	◆	0.88 (0.81 to 0.96)	◆
Fatal or non fatal stroke	0.83 (0.76 to 0.92)	◆	0.81 (0.74 to 0.90)	◆
All-cause mortality	0.88 (0.82 to 0.94)	◆	0.87 (0.81 to 0.94)	◆
Incident HHF	0.89 (0.82 to 0.98)	◆	0.88 (0.79 to 0.98)	◆
Kidney composite (+ macroalbuminuria)	0.79 (0.73 to 0.87)	◆	0.78 (0.71 to 0.87)	◆
Worsening kidney function (eGFR)	0.86 (0.72 to 1.02)	◆	0.82 (0.69 to 0.98)	◆
	0.3 0.6 1.2		0.3 0.6 1.2	

****eGFR data are median (interquartile range) for SUSTAIN 6 and EXSCEL**
ASCVD, atherosclerotic cardiovascular disease; BMI, body mass index; CV, cardiovascular; CVOT, cardiovascular outcomes trial; ER extended release, EOT, end of treatment , eGFR, estimated glomerular filtration rate; FDA, food and drug administration; GLP-1RA, glucagon like peptide 1 receptor agonist; HHF, hospitalization due to heart failure; I, interval; HbA_{1c}, glycated hemoglobin; HR, hazard ratio; MACE, major adverse cardiovascular events; MI, myocardial infarction; NR, not reported; SGLT2i, Sodium-glucose cotransporter 2 inhibitors; SoC, standard of care; T2D, type 2 diabetes
1. Sattar N et al. Lancet Diabetes Endocrinol. 2021; S2213-8587(21)00203-5, 2. <https://clinicaltrials.gov/study/NCT03914326>

